

Panadime™

Description:

Panadime™ is designed for nutritional support of the pancreas by providing enzyme-rich non-processed food and specific nutritional support for pancreatic exocrine and endocrine function.

Indications: Pancreas

Clinical Biochemistry: See introduction.

Precautions, Adverse Reactions, Contraindications:

Digestive upset or diarrhea may result if over consumed or not gradually introduced. Do not use for pregnant or nursing animals.

Note: In acute pancreatitis the stimulation of pancreatic exocrine secretion by high protein diets may promote the release of proteases within pancreatic tissue aggravating the inflammatory response. Fasting and short term low (but high quality) protein diets such as Vegan™ or other Wysong Dry Diets combined with Panadime™ is indicated until biochemical markers are within normal range.

Clinical Biochemistry: See introduction.

Ingredients:

Beef, Beef Liver, Chicken, Chicken Liver, Ground Beef and Chicken Bone, Fish Oil, Coconut Oil, Pepsin, Ox Bile, Pancreatin, Dried Pancreas, Garlic, Lecithin, Dandelion Root, Grape Seed Extract, Milk Calcium, Plums, Lactoferrin, Lactoperoxidase, Barley Grass, Wheat Grass, Desiccated Sea Plankton, Artichoke, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus plantarum Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Dried Lactobacillus casei Fermentation Product, Dried Lactobacillus lactis Fermentation Product, Dried Saccharomyces cerevisiae Fermentation Product, Dried Aspergillus oryzae Fermentation Product, Dried Aspergillus niger Fermentation Product, Phytase, Natural Extractives of Rosemary, Natural Extractives of Sage, Choline Chloride, Ascorbic Acid, Zinc Proteinate, Iron Proteinate, Vitamin E Supplement, Niacin Supplement, Manganese Proteinate, Calcium Pantothenate, Thiamine Mononitrate, Copper Proteinate, Pyridoxine Hydrochloride, Riboflavin Supplement, Vitamin A Acetate, Folic Acid, Biotin, Vitamin B₁₂ Supplement, Vitamin D₃ Supplement.

Analysis: Protein 50.0%, Fat 28%, Fiber 2.8%, Moisture 3.0%

References:

Arivazhagan S, et al. Modulatory effects of garlic and neem leaf extracts on N-methyl-N-nitro-N-nitrosoguanidine (MNGG)-induced oxidative stress in Wistar rats. *Cell Biochem Funct.* 18(1):17-21. 2000.; Bagchi D, et al. Free radicals and grape seed proanthocyanidin extract: importance in human health and disease prevention. *Toxicology.* 148(2-3):187-97. 2000.; Bagchi D, et al. Protective effects of grape seed proanthocyanidins and selected antioxidants against TPA-induced hepatic and brain lipid peroxidation and DNA fragmentation, and peritoneal macrophage activation in mice. *Gen Pharmacol.* 30:771-6. 1998.; Belinky PA, et al. Structural aspects of the inhibitory effect of glabridin on LDL oxidation. *Free Radic Biol Med.* 24(9):1419-29. 1998.; Buchman AL, et al. Lecithin increases plasma free choline and decreases hepatic stenosis in long-term total parenteral nutrition patients. *Gastroenterology.* 102:1363-70. 1992.; Duff G, et al. Lecithin is known to increase the capacity of the bile to solubilize cholesterol. *Am J Med.* 11:92. 1951.; Exogenous enzymes in monogastric nutrition – their current value and future benefits. *Animal Feed Science and Technology.* 86(1-2). 2000.; Francischetti IM, et al. Identification of glycyrrhizin as a thrombin inhibitor. *Biochem Biophys Res Commun.* 235(1):259-63. 1997.; Gabetta B, et al. Characterization of proanthocyanidins from grape seeds. *Fitoterapia.* 71:162-75. 2000.; Germain E, et al. In vivo metabolism of diallyl disulphide in the rat: identification of two new metabolites. *Xenobiotica.* 32(12):1127-38. 2002.; Green M, et al. Inhibition of DES-induced DNA adducts by diallyl sulfide: implications in liver cancer prevention. *Oncol Rep.* 10(3):767-71. 2003.; Hattori A, et al. Protective effect of ajoene on acetaminophen-induced hepatic injury in mice. *Biosci Biotechnol Biochem.* 65(11):2555-7. 2001.; Howell E. **Food enzymes for health and longevity.** Connecticut, Omagond Press. 1980.; Hu C, et al. Antioxidant, prooxidant, and cytotoxic activities of solvent-fractionated dandelion (*Taraxacum officinale*) flower extracts in vitro. *J Agric Food Chem.* 51(1):301-10. 2003.; Jaroni D, et al. The effect of dietary wheat middlings and enzyme supplementation II: Applied nutrient digestibility, digestive tract size, gut viscosity, and gut morphology in strains of leghorn hens. *Poult Sci.* 78(12):1664-74. 1999.; Joshi SS, et al. The cellular and molecular basis of health benefits of grape seed proanthocyanidin extract. *Curr Pharm Biotechnol.* 2(2):187-200. 2001.; Kashiwada Y, et al. Sesquiterpene glucosides from anti-leukotriene B4 release fraction of *Taraxacum officinale*. *J Asian Nat Prod Res.* 3(3):191-7. 2001.; Kim HM, et al. *Taraxacum officinale* inhibits tumor necrosis factor- α production from rat astrocytes. *Immunopharmacol Immunotoxicol.* 22(3):519-30. 2000.; Kisiel W, et al. Further sesquiterpenoids and phenolics from *Taraxacum officinale*. *Fitoterapia.* 71(3):269-73. 2000.; Knowles RP, et al. Clinical impressions of the use of an enzyme additive in large and small animals. *VM/SAC.* 74(12):1733. 1979.; Lau B. **Garlic for health.** Lotus Light Publications, Wilmot, WI. 32-3. 1988.; Layer P, et al. Pancreatic enzymes: Secretion and luminal nutrient digestion in health and disease. *J Clin Gastroenterol.* 28(1):3-10. 1999.; Lewis GE, et al. Effect of direct-fed fibrolytic enzymes on the digestive characteristics of a forage-based diet fed to beef steers. *J Anim Sci.* 74(12):3020-8. 1996.; Matsui T, et al. Efficacy of phytase in improving phosphorus bioavailability in a corn-soybean meal-based diet for growing pigs. *J Anim Sci.* 78:94-9. 2000.; Mavromichalis I, et al. Enzyme supplementation and particle size of wheat in diets for nursery and finishing pigs. *J Anim Sci.* 78:3086-95. 2000.; Polichetti E, et al. Cholesterol-lowering effect of soybean lecithin in normolipidaemic rats by stimulation of biliary lipid secretion. *Br J Nutr.* 75(3):471-8. 1996.; Racz-Kotilla E, et al. The action of *Taraxacum officinale* extracts on the body weight and diureses of laboratory animals. *Planta Medica.* 26:212-7. 1974.; Richardson T. **Enzymes: Principles of food science.** Part 1. Ed or Fennema, New York: Marcel Dekker, Inc. 1976.; Susnik F. The present state of knowledge about the medicinal plant *Taraxacum officinale* Weber. *Med Razgl.* 21:323-8. 1982.; Terrell SS, et al. On trial: An enzyme-producing food supplement for dogs. *Vet Med.* 79(11):1367. 1984.; Thomas M, et al. Diallyl disulfide increases rat h-ferritin, L-ferritin and transferrin receptor genes in vitro in hepatic cells and in vivo in liver. *J Nutr.* 132(12):3638-41. 2002.; Tompkins R, et al. Relationship of biliary phospholipid and cholesterol concentrations to the occurrence of human gallstones. *Ann Surg.* 172(6):936-45. 1970.; Tuzhilin S, et al. The treatment of patients with gallstones by lecithin. *Am J Gastroenterol.* 65:231. 1976.; Watson JP, et al. Case report: oral antioxidant therapy for the treatment of primary biliary cirrhosis: a pilot study. *Journal of Gastroenterology and Hepatology.* 10:1034-40. 1999.; Williams CA, et al. Flavonoids, cinnamic acids and coumarins from the different tissues and medicinal preparations of *Taraxacum officinale*. *Phytochemistry.* 42(1):121-7. 1996.; Yang CS, et al. Mechanisms of inhibition of chemical toxicity and carcinogenesis by diallyl sulfide (DAS) and related compounds from garlic. *J Nutr.* 131(3s):1041S-5S. 2001.; Yang WZ, et al. Effects of an enzyme feed additive on extent of digestion and milk production in lactating dairy cows. *J Dairy Sci.* 82(2):391-403. 1999.; Zhao J, et al. Anti-tumor-promoting activity of a polyphenolic fraction isolated from grape seed in the mouse skin two-stage initiation-promotion protocol and identification of procyanidin B5-3'-gallate as the most effective antioxidant constituent. *Carcinogenesis.* 20:1737-45. 1999.