

Vermiden™

Description:

Vermiden™ is designed to provide nutritional support for animals suffering from intestinal parasites, which jeopardize the health of the G/I lining and compromise nutritional status.

Indications: Parasites

Clinical Biochemistry: See introduction.

Precautions, Adverse Reactions, Contraindications:

Digestive upset or diarrhea may result if over consumed or not gradually introduced. Do not use for pregnant or nursing animals.

Ingredients:

Beef, Beef Liver, Chicken, Chicken Liver, Ground Beef and Chicken Bone, Fish Oil, Coconut Oil, Desiccated Sea Plankton, Garlic, Black Walnut, Milk Calcium, Plums, Lactoferrin, Lactoperoxidase, Barley Grass, Wheat Grass, Artichoke, Amylase, Protease, Lipase, Cellulase, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus plantarum Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Dried Lactobacillus casei Fermentation Product, Dried Lactobacillus lactis Fermentation Product, Dried Saccharomyces cerevisiae Fermentation Product, Dried Aspergillus oryzae Fermentation Product, Dried Aspergillus niger Fermentation Product, Phytase, Natural Extractives of Rosemary, Natural Extractives of Sage, Choline Chloride, Ascorbic Acid, Zinc Proteinate, Iron Proteinate, Vitamin E Supplement, Niacin Supplement, Manganese Proteinate, Calcium Pantothenate, Thiamine Mononitrate, Copper Proteinate, Pyridoxine Hydrochloride, Riboflavin Supplement, Vitamin A Acetate, Folic Acid, Biotin, Vitamin B₁₂ Supplement, Vitamin D₃ Supplement.

Analysis: Protein 50.0%, Fat 28%, Fiber 2.8%, Moisture 3.0%

References:

Adetumbi M, et al. *Allium sativum* (garlic) – a natural antibiotic. *Medical Hypoth.* 12:227-37. 1983.; Anesini C, et al. Screening of plants used in Argentine folk medicine for antimicrobial activity. *Journal of Ethnopharmacology.* 39:119-28. 1993.; Ankri S, et al. Antimicrobial properties of allicin from garlic. *Microbes Infect.* 1(2):125-9. 1999.; Barr SC, et al. Giardiasis of dogs and cats. *Comp Cont Edu Pract Vet.* 16:603-14. 1994.; Cellini L, et al. Inhibition of *Helicobacter pylori* by garlic extract (*Allium sativum*). *FEMS Immunology and Medical Microbiology.* 13:273-7. 1996.; Chowdhury AKZ, et al. Efficacy of aqueous extract of garlic and allicin in experimental shigellosis in rabbits. *Indian J of Med Res.* 93:33-6. 1991.; Cline S, et al. Determination of methyl 2-benzimidazolylcarbamate in black walnut fruit. *J Agric Food Chem.* 29(5):1087-8. 1981.; Collins GH. A limited survey of gastro-intestinal helminths of dogs and cats. *NZ Vet J.* 21:175-6. 1973.; Curreli N, et al. Effects of plant-derived naphthoquinones on the growth of *Pleurotus sajor-caju* and degradation of the compounds by fungal cultures. *J Basic Microbiol.* 41(5):253-9. 2001.; Davis LE, et al. In vitro synergism of concentrated *Allium sativum* extract and amphotericin B against *Cryptococcus neoformans*. *Planta Medica.* 60:546-9. 1994.; Fetterer RH, et al. Effects of plumbagin on development of the parasitic nematodes *Haemonchus contortus* and *Ascaris suum*. *Comp Biochem Physiol C.* 100(3):539-42. 1991.; Foster S, et al. **Black walnut in medicinal plants.** Houghton Mifflin Co. New York, NY. 276. 1990.; Galey FD, et al. Black walnut (*Juglans nigra*) toxicosis: a model for equine laminitis. *J Comp Pathol.* 104(3):313-26. 1991.; Gallwitz H, et al. Ajoene is an inhibitor and subversive substrate of human glutathione reductase and *Trypanosoma cruzi* trypanothione reductase: crystallographic, kinetic, and spectroscopic studies. *J Med Chem.* 42(3):364-72. 1999.; Hasslinger MA. *Ollulanus tricuspis*, the stomach worm of the cat. *Feline Pract.* 14:22-35. 1984.; Hennig L, et al. Selective inactivation of parvalin-like peptidyl-prolyl cis/trans isomerases by juglone. *Biochemistry.* 37(17):5953-60. 1998.; Hoshino T, et al. Effects of garlic preparations on the gastrointestinal mucosa. *J Nutr.* 131:1109S-1113S. 2001.; Kamei H, et al. Inhibition of cell growth in culture by quinones. *Cancer Biother Radiopharm.* 13(3):185-8. 1998.; Lawson LD, et al. Characterization of the formation of allicin and other thiosulfonates from garlic. *Planta Medica.* 58:345-50. 1992.; Lawson LD, et al. Pre-hepatic fate of the organosulfur compounds derived from garlic (*Allium sativum*). *Planta Medica.* 59:A688. 1993.; Lee KW, et al. Hematologic changes associated with the appearance of eocentrocytes after intragastric administration of garlic extract to dogs. *Amer J Vet Res.* 61(11):1446-50. 2000.; Lun, et al. Antiparasitic activity of diallyl trisulfide (dasuansu) on human and animal pathogenic protozoa (*Trypanosoma* sp., *Entamoeba histolytica* and *Giardia lamblia*) in vitro. *Ann Soc Belge Med Trop.* 74:51-9. 1994.; Mahoney, et al. Regulation of aflatoxin production by naphthoquinones of walnut (*Juglans regia*). *J Agric Food Chem.* 48(9):4418-21. 2000.; Munday R, et al. Induction of quinone reductase and glutathione transferase in rat tissues by juglone and plumbagin. *Planta Med.* 66(5):399-402. 2000.; Nok AJ, et al. *Allium sativum*-induced death of African trypanosomes. *Parasitology Res.* 82:634-7. 1996.; Singh KV, et al. Activity on multiple resistant bacteria of garlic (*Allium sativum*) extract. *Fitoterapia.* LV(5):313-5. 1984.; Soffer SA, et al. Evaluation of the antiparasitic effect of aqueous garlic (*Allium sativum*) extract in *Hymenolepis nana* and giardiasis. *J Egypt Soc Parasit.* 21(2):497-502. 1991.; Sugie S, et al. Inhibitory effects of plumbagin and juglone on azoxymethane-induced intestinal carcinogenesis in rats. *Cancer Lett.* 127(1-2):177-83.; Weber ND, et al. In vitro virucidal effects of *Allium sativum* (garlic) extract and compounds. *Planta Medica.* 58:417-23. 1992.